



UNIVERSITY OF OSLO

# Petroleum potential and thermal maturity of Palaeozoic sediments from the North Sea region

Jon H. Pedersen<sup>1</sup>, Dr. Dag A. Karlsen<sup>1</sup>, Dr. Jan E. Lie<sup>2</sup> and Harald Brunstad<sup>2</sup>

<sup>1</sup>Department of Geosciences, University of Oslo, P.O. Box 1047 Blindern, N-0316 Oslo, Norway  
<sup>2</sup>RWE Dea Norge AS, P.O. Box 243 Skøyen, N-0213 Oslo, Norway



## Introduction

The dominating oil and gas source rocks in the northern North Sea are of Mesozoic age. Especially important are the Upper Jurassic marine clays of the Kimmeridge Clay Formation, deposited in the Jurassic Moray Firth Basin and Central and Viking Graben rift structures.

There are however Palaeozoic sediments with possible petroleum potential in both Norwegian and Danish wells in the northern North Sea. We also know that source rocks of Cambro-Silurian, Devonian, Carboniferous and Permian age are found in areas fringing the North Sea, i.e. in Norway, Denmark, Sweden, Poland, Germany, the Netherlands, in the UK and on East Greenland.

In the Skagerrak Sea and northern North Sea, thick sequences of marine sediments most likely accumulated in Cambro-Silurian times. Additionally, shales, coals and mudstones of Devonian to Permian age may have been deposited in half-grabens and N-S trending rift basins underlying the Upper Permian Zechstein evaporites.

In this study, we investigate the kerogen types, petroleum potential and thermal maturity of some of these Palaeozoic sediments, and look for traces of Palaeozoic petroleum in reservoir rocks in the North Sea. In the northeastern North Sea and Norwegian-Danish Basin (NDB), Palaeozoic source rocks may be a crucial part of plays capped by thick Upper Permian evaporite layers.




Fig. 1. Some occurrences of Palaeozoic source rocks in the North West Europe.

## Conceptual stratigraphy

Em	Period	Age	Group	Rock unit	Source	Reservoir	Cap	
Cenozoic	Quaternary	1.1-0	Holocene	Bergen				
				Stavanger				
				Leik				
	Tertiary	23.2-0	Pliocene	Nordland				
				Hordaland				
				Rogaland				
	Mesozoic	Cretaceous	Upper	Cromer Knoll	Stige			
					Stige			
					Stige			
		Jurassic	Lower	Boknfjord	Stige			
Stige								
Stige								
Triassic		Middle	Heron	Stige				
				Stige				
				Stige				
Palaeozoic		Permian	Upper	Zechstein	Stige			
	Stige							
	Carboniferous	Upper	Rotliegende	Stige				
				Stige				
	Devonian	Upper	Boknfjord	Stige				
				Stige				
	Silurian	Upper	Stige	Stige				
				Stige				
	Ordovician	Upper	Stige	Stige				
				Stige				
Cambrian	Middle	Stige	Stige					
			Stige					

Fig. 4. Conceptual stratigraphy in the northern North Sea and the Norwegian-Danish Basin, with possible source, reservoir and cap rocks indicated.




Fig. 2. The study area and well data base.




Fig. 3. The concepts of a possible Palaeozoic petroleum system in the Norwegian-Danish Basin. Possible source rocks are Lower Palaeozoic marine shales, Carboniferous coals and mudstones and Permian marine shales and mudstones.

## Petroleum Potential




Fig. 5.  $T_{max}$  – Hydrogen Index (HI) plot for Palaeozoic samples with TOC over 1 wt%. Cambrian-Ordovician and Permian samples are both oil and gas prone, while Devonian-Carboniferous samples are in general gas prone.

Fig. 6. Total Organic Carbon (TOC) plotted against remaining petroleum potential ( $S_2$ ) of the sediment samples. Carboniferous coals and low-mature onshore samples have the highest  $S_2$  values. Devonian and Permian samples have low  $S_2$  values, while low-mature Cambrian-Ordovician shales have high  $S_2$  values.

## Thermal maturity




Fig. 7. A vitrinite reflectivity versus depth plot. Burial and maturity of the samples increases from the western parts (UK) to the eastern parts of the North Sea (Norway and Denmark). The majority of samples are in the oil generating zone (the oil window).




Fig. 8. Tentative burial curves for the Palaeozoic to present in the Norwegian-Danish Basin and the Skagerrak Sea area. Major subsidence episodes occurred during the Silurian-Devonian Caledonian orogenic event, and as post-rift thermal subsidence in the Triassic subsequent of Permian rifting and volcanism. The Skagerrak Sea area was uplifted and eroded in Neogene times.

## Conclusions

Palaeozoic sediments of Cambrian-Ordovician, Devonian-Carboniferous and Permian age are found in wells in the northern North Sea and Skagerrak Sea

Palaeozoic mudstones and shales from the northern North Sea have TOC values in the range of 0.1-14.4 wt%.

Upper Palaeozoic coals from the northern North Sea have TOC values in the range of 17-80 wt%.

Cambrian-Ordovician and Permian samples are oil and gas prone.

Devonian and Carboniferous samples from the North Sea are gas prone.

The Palaeozoic northern North Sea samples in this study have thermal maturities within the oil window ( $\%Ro = 0.3-1.3$ ).

Burial, and hence maturity, of the Palaeozoic samples increases from west to east in the northern North Sea.

## Acknowledgments

For supplying samples:  
 NPD, Stavanger, Norway  
 GEUS, Copenhagen, Denmark  
 SGU, Uppsala, Sweden  
 DTI Core Store, Edinburgh, Scotland

For funding:  
 RWE Dea Norge AS  
 For analytical help:  
 UIB, Bergen, Norway

## References

Anderson, A., Dalmen, B., Gee, D. G. & Snel, S. (1985) The Scandinavian Alum Shales. *Sources Geologicae Universitatis, Avhandlingar og uppsatser* 1, 64, 56, 1-50

Allen, T. R. (1992) The Devonian lacustrine sediments of Shetland, Scotland: implications for thermal maturity. *Journal of the Geological Society*, London, 147, 145-151

Bird, B. M. (1990) Carboniferous, Devonian, C. W. (ed) *Petroleum Geology of the North Sea* 4th Edition, Blackwell Science Ltd, 104-136

Bjorck, S., Larter, S. & Horsfield, B. (1992) The unusual source potential of the Cambrian Alum Shale in Scandinavia as determined by quantitative pyrolysis methods. *Spencer, A. M. (ed) Generation, accumulation and production of kerogen hydrocarbons: C. Special Publications of the European Association of Petroleum Geoscientists No. 2*, Springer-Verlag Berlin Heidelberg, 103-110

Bjorck, M., Hal, P. B., Laberg, R., McDermott, J. A. & Mils, N. (1995) Hydrocarbons from non-marine source rocks. *Organic Geochemistry*, 13, 1-3, Pergamon Press Ltd, 221-244

Borghesen, A. P., Kevins, S. V., Margolin, S. S. & Pomeroyev (1992) Geology and hydrocarbon prospects of the Paleozoic in the Baltic region. *Parker, J. R. (ed) Petroleum Geology of Northwest Europe: Proceedings of the 4th Conference, The Geological Society, London*, 651-656

Buchner, B., Claxton, J. & Thomson, E. (1986) Carbon isotope composition of Lower Paleozoic kerogens: Effects of maturation. *Organic Geochemistry*, 10, Pergamon Journals Ltd, 124-134

Ruggie, T., Briggs, J. E., Leith, D. A., Margolin, S. S., Wilson, N. M. & Lohm, R. L. (2002) Upper Permian as a new play model on the mid-Norwegian continental shelf: Investigated by shallow stratigraphic drilling. *The American Association of Petroleum Geologists Bulletin*, 82, 5 (May 2002), 101-127

Cameron, I. B. & Stephenson, D. (1985) British Regional Geology, The Midland Valley of Scotland (2nd edition). Natural Environment Research Council, Institute of Geological Sciences, Her Majesty's Stationery Office, London

Christiansen, P. E., Olsen, H., Pedersen, J. & Skjold, S. (1988) Organic geochemistry of Upper Palaeozoic lacustrine shales in the East Greenland Basin. *Organic Geochemistry*, 18, 1-3, Pergamon Press Ltd, 287-294

Comber, C. (1995) Source rocks and hydrocarbons of the North Sea. *Claxton, J. R. (ed) Petroleum Geology of the North Sea 4th Edition*, Blackwell Science Ltd, 376-462

Doris, A. A. (1956) Coal blend and varietal hydrocarbon in Norway. *Norsk Geologisk Tidsskrift*, 34, 250-260

Esteban, G., Manning, D. A. C. & Abbott, G. D. (1992) An organic geochemical study of bitumens and their potential source rocks from the South Permian, Ordovician, Central England. *Organic Geochemistry*, 20, 5, 579-596

Fallick, B. & Tyson, R. V. (1982) Organic geochemistry of Upper Palaeozoic lacustrine shales in the East Greenland Basin. *Organic Geochemistry*, 5, 1-3, Pergamon Press Ltd, 287-294

Garand, J., Shabbar, T. J., Blöcher, J. S., Sultan, M. & Bassett, M. G. (1992) Permian-Carboniferous and other plays: Their historical development and future potential. *Parker, J. R. (ed) Petroleum Geology of Northwest Europe: Proceedings of the 4th Conference, The Geological Society, London*, 441-460

Halsrud, M. (1991) Large podmorans, gas-charged sediments and possible clay dikes in the Skagerrak. *Marine and Petroleum Geology*, 8, 3, Butterworth-Heinemann Ltd, 311-316

Wang, H. & Meyer, T. (1985) Lacustrine organic facies: A kinematic study using multivariate statistical analysis. *Organic Geochemistry*, 8, 2, 191-210

Karlsen, D. A., Nordviken, T., Larter, S. & Björck, K. (1992) Hydrocarbon composition of authigenic inclusions: Application to elucidation of petroleum reservoir filling history. *Geochimica et Cosmochimica Acta*, 57, 3641-3659

Liibe, R., Leythaeuser, D., Radde, M. & Schuler, R. G. (1990) Petroleum generation and migration in coal seams of the Carboniferous Ruhr Basin, northwestern Germany. *Organic Geochemistry*, 18, 1-3, Pergamon Press Ltd, 247-258

Olaussen, S. (1981) Marine incursion in Upper Palaeozoic sedimentary rocks of the Oslo Region, Southern Norway. *Geological Magazine*, 118, 2, 261-286

Peters, K. E., Moldauer, S. M., Orndorff, B. & Cernomach, G. J. (1989) Origin of Basins Oil by Co-Generation from Devonian and Middle Jurassic Source Rocks, Inner Shetland Firth, United Kingdom. *The American Association of Petroleum Geologists Bulletin*, 73, 4, 454-471

Skjold, S., Christiansen, P. E. & Pedersen, J. (1992) Carboniferous lacustrine shale in East Greenland – additional source rock in the northern North Atlantic. *Katz, B. J. (ed) Lacustrine basin evolution: case studies and modern analogues*. *American Association of Petroleum Geologists*, 50, 277-286

Sorenson, S. & Martinius, B. W. (1973) A paleogeographic reconstruction of the Rotliegende deposits in the northeastern Permian Basin. *Brooks, J. & Galloway, K. (eds) Petroleum Geology of North West Europe*, Graham & Trotman, 497-503

## Introduction

The Palaeozoic successions in the northern North Sea are not as well mapped and understood as younger strata in the North Sea sedimentary basins. This is due to great burial depths, but also because the Palaeozoic has been regarded as of low economic interest in this area.

In this project we investigate Palaeozoic sediments collected from wells and outcrops in the North Sea and adjacent areas. In specific, we focus on Cambro – Silurian, Carboniferous and Permian sediments, which may have petroleum source rock properties.

We hope that this study will increase our understanding of the distribution, organic facies and thermal maturity of Palaeozoic sediments in the northern North Sea. We also believe that this work may help us to better evaluate the significance of Palaeozoic petroleum systems in the northern North Sea and Norwegian Danish Basin.




Fig.1. Study area and samples selected for pyrolysis analysis.

## Methodology

The sample set contains outcrop samples, core chips and cuttings from selected locations and wells in the North Sea and adjacent areas. The samples represent Cambrian, Ordovician, Carboniferous and Permian sediments of both marine and continental origin.

The samples underwent the following analytical procedures:

- Soxhlet extraction of sediments with DCM:MeOH (97:3)
- GC-FID and GC-MS analysis of solvent extracts
- Kerogen up-concentration from sediment samples using HF and HCl acid
- PY-GC of kerogen concentrates, (open pyrolysis, 300°C-600°C in 10 minutes in a helium atmosphere)

## Results




Fig.2. Chromatograms of pyrolysis products (PY-GC) and solvent extracts (GC-FID) of Palaeozoic samples. The samples are of low to moderate maturity. Identifiable compounds on the pyrograms are pairs and aromatic compounds like benzene, toluene and naphthalene. The isoprenoids pristane and phytane are not generated during pyrolysis. The products from the marine Alum shale are dominated by low waxiness and high amounts of aromatic compounds. The organic rich Carboniferous Midland Valley lacustrine shale generates a high-wax product with an odd-even predominated n-alkane distribution, while the marine/hypersaline Permian Kupferschiefer gives a product with a typical marine n-alkane envelope. Note however the enrichments of C<sub>13</sub>-C<sub>16</sub> n-alkanes on the Kupferschiefer GC-FID chromatogram.




Fig.3. A plot of phytane/n-C<sub>18</sub> versus pristane/n-C<sub>17</sub> is used here to evaluate organic facies, thermal maturity and degree of biodegradation of oils and extracts. Cambrian-Ordovician and Permian samples have a marine affinity, resembling Jurassic oils and extracts. Extracts from Devonian/Carboniferous sediments points to a lacustrine or transitional environment (Modified from Shanmugam, 1985).




Fig.4. A ternary plot of the percentages of the three main groups of n-alkanes in the pyrolysates. Coals and organic lean mudstones generates gas only, while algae rich coals and marine shales produce a low-wax oil. Two organic rich Carboniferous lacustrine shales generates a high wax paraffinic oil. (Adapted from Horsfield, 1989).




Fig.5. A ternary plot of an aromatic compound (toluene), an n-alkane component (C<sub>9</sub>+C<sub>22</sub>) and a sulphur-compound (2,5-dimethylthiophene) identified in the pyrolysates. Organic lean mudstones have high percentages of aromatic compounds, while the majority of samples, both coals and shales classifies as intermediate, but biased towards the aromatic corner. (Adapted from Horsfield, 1989).




Fig.6. The maturity of extracts and oils expressed by the aromatic compound methylphenanthrene (Kvalheim et al., 1987). The samples are in general of oil generating maturity. Shallow samples are from outcrops or onshore wells.

## Conclusions

Carboniferous samples (coals and mudstones) from the North Sea generates mainly gas under open pyrolysis.

Cambrian-Ordovician and Permian samples (shales) generate both gas and oil products under open pyrolysis.

The samples are within the oil window in terms of thermal maturity.

GC-FID data suggest Cambrian-Ordovician and Permian samples to have a marine origin.

GC-FID data suggest Devonian-Carboniferous samples to have a lacustrine or deltaic origin.

Palaeozoic coals, shales and mudstones in the North Sea region have probably generated petroleum.

## Acknowledgments

For supplying samples:  
NPD, Stavanger, Norway  
GEUS, Copenhagen, Denmark  
SGU, Uppsala, Sweden  
DTI Core Store, Edinburgh, Scotland

For funding:  
RWE Dea Norge AS

## References

Allen, J., Bjeroy, M. & Douglas, A. C. (1975) A geochemical study of the exinite group maceral alginite, selected from three Permian-Carboniferous torbanites. *Diagenesis*, **1**, 631-674.

Bruland, S., Larsen, S. & Horsfield, B. (1992) The unusual source potential of the Cambrian Alum Shale in Scandinavia as determined by quantitative pyrolytic methods. *Spectrochim. Acta B*, **16**, 1055-1062.

Burford, A. C., Karlsen, D. A., Baskin, K., Lu, T., Skjerve, L., Bernthansen, H. H. & Kihlström, J. E. (2000) Reservoir characterization by a combined micro-analysis - micro thin-layer chromatography (GC-MS) method. A calibration study with samples from the Norwegian North Sea. *Journal of Petroleum Geology*, **23**, 227-244.

Burford, A. C. & Hansen, M. (2000) Oriskanyite I oils - an specialised fra Ordovician and Kinshofen. *Surv.*, **3**, 1-7.

Christiansen, F. G., Larsen, H. C., Marzouk, C., Plassard, S. & Steinemærk, L. (1992) Late Palaeozoic plays in East Greenland. *Petroleum*, **1**, 1-10.

Esteban, G. (1994) High-temperature programmed pyrolysis of Palaeozoic source rocks from Northern Germany and adjacent areas and its thermodynamic constraints. *Organic Geochemistry*, **24**, 1011-1025.

Esteban, G., Manning, D. A. C. & Abbott, C. D. (1992) An organic geochemical study of bitumens and their potential source rocks from the South Pennine Orefield, Central England. *Organic Geochemistry*, **20**, 579-590.

Horsfield, B., Dixon, U. & Leister, J. (1989) The microscale simulation simulation of maturation: Outline of a new technique and its potential applications. *Geochimica et Cosmochimica Acta*, **53**, 317-374.

Kim, J. (1990) Application of biomarker oxidation-esterification micro-spectrometry for characterization of high hydrocarbon fluid inclusion. *Organic Geochemistry*, **23**, 111-112.

Kishinoue, O., M. Chino, A. A., Tsuboi, N. & Sugimoto, A. (1987) Maturity determination of organic matter in shale using the methylphenanthrene distribution. *Geochimica et Cosmochimica Acta*, **51**, 1863-1880.

Lie, J. E. (1996) Deep seismic reflection mapping of crustal and upper mantle structures beneath the Saggeren Sea, SW Scandinavia. *Dissertation for the degree Doctor Scientiarum*. Department of Geology, Faculty of Mathematics and Natural Sciences, University of Oslo.

Pedersen, J. H. (1982) Hydrocarbon source rocks, source rock and migration in the Orkney Basin. *Scottish Journal of Geology*, **28**, 327-336.

Pedersen, J. H. (2000) Algalite Oils and Condensates of the North Atlantic margin of Pangaea. *Organic Geochemical Society*. *Field School*. *University of Oslo*, Norway.

Steinemærk, L. (2000) Late Palaeozoic evolution of the North Atlantic margin of Pangaea. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **161**, 91-126.

Thompson, A. & Leggett, J. K. (1987) Stratigraphic distribution and palaeo-geochemical significance of European early Palaeozoic organic-rich sandstones. *Bull. J. & Flett, A. J. (eds) Marine Petroleum Source Rocks*, Geological Society Special Publication **26**, 323-347.